Midterm Review Session problems

October 6, 2025

Exercise 1 (Filling a contingency table from conditionals): You are told that (X, Y) is a pair of random variables where $X \in \{0, 1\}$ and $Y \in \{0, 1, 2\}$. The joint distribution is not given, but you are told the following information:

$$P(Y = 0) = 1/2$$
, $P(Y = 1) = 1/4$, $P(Y = 2) = 1/4$

and the conditional probabilities of X given Y are

$$P(X = 1 | Y = 0) = 1/2$$
, $P(X = 1 | Y = 1) = 1/4$, $P(X = 1 | Y = 2) = 1/4$

Fill in the contingency table for the joint distribution P(X = x, Y = y):

$$X = 0$$
 $Y = 1$ $Y = 2$ $X = 1$ $X = 1$

Exercise 2 (Covariance of two Bernoulli random variables): Let (X, Y) be a pair of Bernoulli random variables with joint distribution given by

$$X = 0$$
 $Y = 0$ $Y = 1$ $X = 0$ 0.3 0.2 $X = 1$ 0.1 0.4

Compute cov(X, Y).

Exercise 3 (Sample distribution): Consider the problem of estimating the mean of a normal random variable $X \sim \text{Normal}(\mu, \sigma^2)$ with known variance σ^2 . You have N iid samples X_1, \ldots, X_N of X where μ is unknown. Write down an estimator whose sample distribution is

$$\hat{\mu}_{\odot} \sim \text{Normal}\left(\mu a + c/N, \frac{a^2 \sigma^2}{N}\right), \quad a, c \neq 0$$

Exercise 4: In class we learned that given N samples X_1, \ldots, X_N from a Normal distribution with unknown mean

$$\hat{\sigma}_1^2 = \frac{1}{N} \sum_{i=1}^N (X_i - \overline{X})^2$$

is a biased estimator of the variance. Explain (without writing out any code) how you could test that this is indeed biased in Python. Specifically, describe what data you would generate and a plot you could make.

Exercise 5 (Translating code to math): What will the following code print?

```
> import numpy as np
> n = 1000
> x = np.random.binomial(3,1/2,size=n)
> y = np.random.normal(1+ x,1,n)
> print(np.mean(y))
```

Exercise 6 (Translating math to code): Consider the model

$$X_1 \sim \mathsf{Bernoulli}(1/2)$$

 $X_2|X_1 \sim \mathsf{Bernoulli}(X_1/2 + 1/4)$
 $X_3|X_2 \sim \mathsf{Normal}(X_1 - X_2, X_1 + 1)$

Write a python code to approximate $P(X_3 < 2|X_2 = 0)$.

Exercise 7 (Recovering regression parameters): Let


$$Y \mid X \sim \text{Normal}(\beta_0 + \beta_1 X, \ \sigma_{\varepsilon}^2),$$
 (1)

Given the marginal mean, expectation and correlation coefficient

$$\mu_X = 2$$
, $\sigma_X^2 = 1$, $\mu_Y = 5$, $\sigma_Y^2 = 4$, $\rho = 1/2$

Find β_1 , β_0 and σ_{ε}^2 .

Exercise 8: The following is a plot from the paper *Lithium deficiency and the onset of Alzheimer's disease* published in Nature. It shows Lithium levels in two groups (NCI = no cognitive impairment, AD = Alzheimer's disease). Why might a linear regression model with the group as the predictor X and Lithium level as the response Y be inappropriate for this data?



Solutions

Exercise 1: Filling a contingency table from conditionals

Using

$$P(X = 1, Y = y) = P(X = 1 | Y = y) P(Y = y),$$
 $P(X = 0, Y = y) = (1 - P(X = 1 | Y = y)) P(Y = y).$ (2)

we get

which sum to 1 as a check.

Exercise 2: Covariance of two Bernoulli random variables

Compute the needed expectations:

$$\mathbb{E}[X] = P(X=1) = 0.1 + 0.4 = 0.5,\tag{3}$$

$$\mathbb{E}[Y] = P(Y=1) = 0.2 + 0.4 = 0.6,\tag{4}$$

$$\mathbb{E}[XY] = 1 \cdot 1 \cdot P(X = 1, Y = 1) = 0.4. \tag{5}$$

Hence,

$$Cov(X,Y) = \mathbb{E}[XY] - \mathbb{E}[X]\mathbb{E}[Y] = 0.4 - (0.5)(0.6) = 0.1.$$
(6)

Exercise 3: Sample distribution

Let $\overline{Y}_N = \frac{1}{N} \sum_{i=1}^N Y_i$. Then

$$\overline{Y}_N \sim \text{Normal}\left(\mu, \frac{\sigma^2}{N}\right).$$
 (7)

Define the estimator

$$\widehat{\mu}_{\widehat{\odot}} = a\overline{Y}_N + \frac{c}{N}. \tag{8}$$

By linear transformation of a Normal,

$$\widehat{\mu}_{\odot} \sim \text{Normal}\left(a\mu + \frac{c}{N}, \frac{a^2\sigma^2}{N}\right),$$
 (9)

which matches the required form.

Exercise 4: Testing bias of $\hat{\sigma}_1^2$ in Python (no code required)

To empirically demonstrate bias of

$$\hat{\sigma}_1^2 = \frac{1}{N} \sum_{i=1}^{N} (X_i - \overline{X})^2, \tag{10}$$

you could:

- 1. Fix a ground-truth Normal model, e.g. $X_i \sim \text{Normal}(\mu_0, \sigma_0^2)$ with known μ_0, σ_0 and a chosen N.
- 2. Repeat for many trials (e.g. 50,000): simulate a size-N = 10 sample, compute $\hat{\sigma}_1^2$ for each trial.
- 3. Plot the sampling distribution (histogram) of $\hat{\sigma}_1^2$ and overlay a vertical line at σ_0^2 . The sample mean

$$\overline{\hat{\sigma}_{1}^{2}} = \frac{1}{T} \sum_{t=1}^{T} \hat{\sigma}_{1,t}^{2} \tag{11}$$

will lie systematically below σ_0^2 , visualizing the negative bias.

Exercise 5: Translating code to math

The model is a linear regression model, using the formula $\mu_Y = \beta_0 + \beta_1 \mu_X$ we get

$$\mathbb{E}[y] = 1 + \frac{3}{2} = \frac{5}{2} \tag{12}$$

Therefore the code will print a number close to 2.5.

Exercise 6: Translating math to code

```
import numpy as np
N = 100000
x1 = np.random.binomial(1, 1/2, size=N)
x2 = np.zeros(N)
x3 = np.zeros(N)
for i in range(N):
    x2[i] = np.random.binomial(1, x1[i]/2 + 1/4)
    x3[i] = np.random.normal(x1[i] - x2[i], np.sqrt(x1[i] + 1))

print(np.mean(x3[x2==0]<2))
# or
print(len(x3[(x2==0) & (x3<2)])/len(x3[x2==0]))</pre>
```

Exercise 7: Recovering regression parameters

Under the model $Y \mid X \sim \text{Normal}(\beta_0 + \beta_1 X, \sigma_{\epsilon}^2)$, we have

$$Cov(X,Y) = \beta_1 Var(X), \tag{13}$$

$$\rho = \frac{\mathsf{Cov}(X, Y)}{\sigma_X \sigma_Y} = \beta_1 \frac{\sigma_X}{\sigma_Y}. \tag{14}$$

Hence

$$\beta_1 = \rho \frac{\sigma_Y}{\sigma_X} = \frac{1}{2} \cdot \frac{2}{1} = 1. \tag{15}$$

Then

$$\beta_0 = \mathbb{E}[Y] - \beta_1 \mathbb{E}[X] = 5 - 1 \cdot 2 = 3,$$
 (16)

and using $Var(Y) = \beta_1^2 Var(X) + \sigma_{\varepsilon}^2$,

$$\sigma_{\varepsilon}^2 = \sigma_Y^2 - \beta_1^2 \sigma_X^2 = 4 - 1^2 \cdot 1 = 3. \tag{17}$$

Exercise 9: Regression appropriateness for the lithium plot

The variance var(Y|X) appears to differ between groups.